Table 4 Broadscale Habitat Affinities of Ranked Taxa on the List of California Bird Species of Special Concern 2002^a

| Taxon | MA^b | WE^c | RI^d | CF ^e | MF^{f} | OW^g | DW^h | SC^{i} | GR^{j} |
|------------------------------------|--------|--------|--------|-----------------|----------------------------|--------|--------|----------|----------|
| Fork-tailed storm-petrel | X | | | | | | | | |
| Ashy storm-petrel | X | | | | | | | | |
| Black storm-petrel | X | | | | | | | | |
| American white pelican | | X | | | | | | | |
| Least bittern | | X | | | | | | | |
| Wood stork | | X | | | | | | | |
| Fulvous whistling-duck | | X | | | | | | | |
| Tule greater white-fronted goose | | X | | | | | | | |
| Brant | | X | | | | | | | |
| Redhead | | X | | | | | | | |
| Harlequin duck | | X | | | | | | | |
| Northern harrier | | X | | | | | | | X |
| Harris's hawk | | | X | | | | | | |
| Prairie falcon | | | | | | | | | X |
| Greater sage-grouse | | | | | | | | X | |
| Mount Pinos blue grouse | | | | X | | | | | |
| Catalina California quail | | | | | | | | X | |
| Yellow rail | | X | | | | | | | |
| Lesser sandhill crane | | X | | | | | | | X |
| Snowy plover (interior population) | | X | | | | | | | 11 |
| Mountain plover | | | | | | | | | X |
| Gull-billed tern | | X | | | | | | | |
| Royal tern | | X | | | | | | | |
| Elegant tern | | X | | | | | | | |
| Black tern | | X | | | | | | | |
| Black skimmer | | X | | | | | | | |
| Xantus's murrelet | X | | | | | | | | |
| Cassin's auklet | X | | | | | | | | |
| Tufted puffin | X | | | | | | | | |
| Burrowing owl | | | | | | | | | X |
| California spotted owl | | | | X | | | | | |
| Long-eared owl | | | X | X | | X | X | | |
| Short-eared owl | | X | 21 | 71 | | ** | 21 | | X |
| Vaux's swift | | 7.1 | | X | | | | | 7.1 |
| Olive-sided flycatcher | | | | X | | | | | |
| Vermilion flycatcher | | | X | 4 1 | | | | | |
| Island loggerhead shrike | | | 21 | | | | | X | |
| Gray vireo | | | | | | | X | X X | |
| Catalina Hutton's vireo | | | | | | X | 11 | Λ | |
| California horned lark | | | | | | Λ | | | v |
| Purple martin | | | X | X | | X | | | X |
| i dipic marun | | | Λ | Λ | | Λ | | | |

| Taxon | MA^b | WE^c | RI^d | CF ^e | MF^{f} | OW^g | \mathbf{DW}^{h} | SC^{i} | GR^{j} |
|--|--------|--------|--------|-----------------|----------------------------|--------|-------------------|----------|----------|
| San Diego cactus wren | | | | | | | | X | |
| Clark's marsh wren | | X | | | | | | | |
| California Swainson's thrush | | | X | | X | | | | |
| Bendire's thrasher | | | | | | | X | X | |
| Crissal thrasher | | | X | | | | | X | |
| Le Conte's thrasher (San Joaquin population) | | | | | | | | X | |
| Lucy's warbler | | | X | | | | X | | |
| Yellow warbler | | | X | | | | | | |
| Sonora yellow warbler | | | X | | | | | | |
| San Francisco common yellowthroat | | X | X | | | | | | |
| Yellow-breasted chat | | | X | | | | | | |
| Summer tanager | | | X | | | | | | |
| San Clemente spotted towhee | | | | | | | | X | |
| Santa Cruz Is. rufous-crowned sparrow | | | | | | | | X | |
| Oregon vesper sparrow | | | | | | | | | X |
| Bryant's savannah sparrow | | X | | | | | | | X |
| Large-billed savannah sparrow | | X | | | | | | X | |
| Grasshopper sparrow | | | | | | | | | X |
| song sparrow ("Modesto" population) | | X | | | | | | | |
| Suisun song sparrow | | X | | | | | | | |
| Samuel's song sparrow | | X | | | | | | | |
| Alameda song sparrow | | X | | | | | | | |
| Channel Island song sparrow | | | | | | | | X | |
| Kern red-winged blackbird | | X | | | | | | | X |
| Tricolored blackbird | | X | | | | | | | X |
| Yellow-headed blackbird | | X | | | | | | | |

^a Species classified on the basis of their primary use of various broad classes of habitats; some species classified as having more than one primary habitat.

b marine (nearshore, offshore, and pelagic waters).

^c wetlands (tidal flats, tidal marsh, freshwater marsh, wet meadows, vemal pools, flooded agricultural fields, and riverine, lacustrine, and estuarine waters). Italics indicates taxa that in their season of concern use primarily coastal or near-coastal (e.g., Salton Sea) saline habitats; all other use primarily interior wetlands except for the San Francisco common yellowthroat and the Bryant's savannah sparrow, which use a combination of estuarine marshes and either freshwater marshes or moist upland grasslands, respectively.

^d riparian forest and woodland. Italics indicate taxa that use primarily desert riparian habitats.

^e coniferous forest.

^f mixed evergreen hardwood forest.

^g oak woodland and oak savannah.

^h desert woodland (Joshua tree, fan palm, mohave yucca, ocotillo, and pinyon-juniper).

ⁱ scrub habitats (chaparral, coastal scrub, desert scrub, and sagebrush scrub).

^j grassland (native grassland, pastureland, grass-like crops, weedy fields, and sparsely-vegetated cultivated fields).